

Theory of Superconductivity, Frühjahrsemester 2026

Blatt 2

Abgabe: 12.03.26, 12:00H (auf adam oder Treppenhaus 4. Stock)

Tutor: Tobias Nadolny, Zi.: 4.48

(1) **Quasi-particle excitations in superconductors** (5 Punkte)

Define operators $\gamma_{\mathbf{k}\uparrow}^\dagger$ and $\gamma_{\mathbf{k}\downarrow}^\dagger$ by

$$\begin{aligned}\gamma_{\mathbf{k}\uparrow}^\dagger &= u_{\mathbf{k}}^* c_{\mathbf{k}\uparrow}^\dagger - v_{\mathbf{k}}^* c_{-\mathbf{k}\downarrow} \\ \gamma_{-\mathbf{k}\downarrow}^\dagger &= u_{\mathbf{k}}^* c_{-\mathbf{k}\downarrow}^\dagger + v_{\mathbf{k}}^* c_{\mathbf{k}\uparrow};\end{aligned}$$

$u_{\mathbf{k}}$ and $v_{\mathbf{k}}$ are complex numbers satisfying $|u_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^2 = 1$ for each momentum \mathbf{k} , and c^\dagger, c are the (standard) electron creation and annihilation operators.

- (a) Prove that the superconducting ground state $|\psi_{\text{BCS}}\rangle = \prod_{\mathbf{k}} (u_{\mathbf{k}} + v_{\mathbf{k}} c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger) |0\rangle$ is the vacuum state of the γ operators, that is,

$$\gamma_{\mathbf{k}\uparrow} |\psi_{\text{BCS}}\rangle = \gamma_{\mathbf{k}\downarrow} |\psi_{\text{BCS}}\rangle = 0.$$

- (b) Obtain explicit expressions of the states created by the γ^\dagger operators, $\gamma_{\mathbf{k}\uparrow}^\dagger |\psi_{\text{BCS}}\rangle$ and $\gamma_{\mathbf{k}\downarrow}^\dagger |\psi_{\text{BCS}}\rangle$, in terms of the electron creation operators $c_{\mathbf{k}\uparrow}^\dagger$ and $c_{\mathbf{k}\downarrow}^\dagger$.

We will see that the states created by $\gamma_{\mathbf{k}\uparrow}^\dagger, \gamma_{\mathbf{k}\downarrow}^\dagger$ are the quasi-particle excitations of wave vector \mathbf{k} and spin \uparrow and \downarrow above the superconducting ground state.

(2) **Average and fluctuations of electron number in the BCS state** (5 Punkte)

- (a) Obtain the average electron number $\bar{N} = \langle \psi_{\text{BCS}} | N | \psi_{\text{BCS}} \rangle$ in the BCS ground state in terms of $v_{\mathbf{k}}$ or $u_{\mathbf{k}}$, where the total electron number operator N has the following second-quantized form:

$$N = \sum_{\mathbf{k}} \left(c_{\mathbf{k}\uparrow}^\dagger c_{\mathbf{k}\uparrow} + c_{\mathbf{k}\downarrow}^\dagger c_{\mathbf{k}\downarrow} \right).$$

Interpret the result in view of the physical meaning of $v_{\mathbf{k}}$ or $u_{\mathbf{k}}$.

Hint: one way (but not the only way) is to rewrite the electron operators in terms of the γ -operators and then use the result of problem 1(a).

- (b) Obtain the fluctuation of the electron number $(\delta N)^2 = \langle \psi_{\text{BCS}} | (N - \bar{N})^2 | \psi_{\text{BCS}} \rangle$ in a similar way as done in (a). How does $\delta N / \bar{N}$ behave in the thermodynamic limit $\bar{N} \rightarrow \infty$?
- (c) (independent of (a) and (b)) Show that

$$|\psi_N\rangle = \left(\sum_{\mathbf{k}} g_{\mathbf{k}} c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger \right)^{\frac{N}{2}} |0\rangle$$

can be obtained by projecting $|\psi_{\text{BCS}}\rangle$ on the subspace of states with particle number N . How is $g_{\mathbf{k}}$ related to $u_{\mathbf{k}}, v_{\mathbf{k}}$?