

Mechanik, Herbstsemester 2020

Blatt 5

Abgabe: 20.10.2020, bitte auf adam in den entsprechenden Ordner.

Ein File pro Abgabe; der Filename sollte Ihren Namen enthalten!

Tutor: Pavel Sekatski, pavel.sekatski@unibas.ch

(1) Lenz-Runge vector

(3 Punkte)

- (a) Show that for the potential $V(r) = -\alpha/r$ there is an additional constant of motion (apart from the energy E and the angular momentum $\mathbf{L}$), the Lenz-Runge vector


$$\mathbf{A} = \dot{\mathbf{x}} \times \mathbf{L} - \alpha \frac{\mathbf{x}}{|\mathbf{x}|}.$$

- (b) What is the direction of $\mathbf{A}$? What happens if the orbit is circular?
 (c) Calculate $\mathbf{x} \cdot \mathbf{A}$ and use this relation to determine the orbit. Compare with the result shown in the lecture. What is the geometrical significance of $|\mathbf{A}|/\alpha$?

(2) Motion of a satellite

(4 Punkte)

We would like to model the motion of the ISS (International Space Station) in the Earth's gravitational field $V(r) = -m\tilde{\alpha}/r$. Consider a toy model of a dumbbell consisting of two point masses $m_1 = m_2 = m$ and a rigid massless connection (length $2a$). We assume that the motion is planar; the center-of-mass motion can be described by polar coordinates r, ϕ , and the orientation of the dumbbell by the angle θ , see figure.



- (a) Write down the Lagrangian L and use it to derive the equations of motion.

Result for L :
$$L = m(\dot{r}^2 + r^2\dot{\phi}^2 + a^2(\dot{\phi} - \dot{\theta})^2) + m\tilde{\alpha} \left(\frac{1}{r_2(r, \theta)} + \frac{1}{r_1(r, \theta)} \right).$$

- (b) Show that there are solutions of the form $r = r_0 = \text{const.}$ and $\dot{\phi} = \text{const.}$, and either $\theta = 0$ or $\theta = \pi/2$. Discuss these two solutions. Give expressions for their periods and expand them in the small parameter a/r_0 . Compare with Kepler's result.

- (c) Bonus points: Integrate the equations of motions numerically for initial conditions $\theta(t=0) \notin \{0, \pi/2\}$ and discuss your results.

(3) **Scattering from a Lennard-Jones potential**

(3 Punkte)

The Lennard-Jones potential is defined by

$$V(r) = 4\epsilon\left(\left(\frac{\sigma}{r}\right)^{12} - \left(\frac{\sigma}{r}\right)^6\right)$$

- (a) Plot the effective potential $V_{\text{eff}}(r)$ for different values of the impact parameter s (i.e., the perpendicular distance between the center of force and the incident velocity). Discuss the qualitatively different cases. What are the characteristic energies?
- (b) Sketch (or calculate numerically) the scattering angle $\Theta(s, E)$ as a function of the impact parameter for the energies found in (a). Give reasons for the shape of the curves and discuss them.

(4) **Bonus problem: Brachistochrone**

(3 Bonus-Punkte)

We return to the situation of problem 1 of Blatt 1: a point mass that slides without friction on a curve $y(x)$ in the xy-plane connecting the two points $r_A = (0, 0)$ and $r_B = (2, -1)$. The mass starts at r_A with velocity 0 and is subject to the Earth's gravitational field that is assumed to be homogeneous and point in the negative y-direction. The total time that the particle needs to reach r_B can be expressed as $T = \int_{r_{Ax}}^{r_{Bx}} dx F(y, y')$ where $F(y, y') = \frac{\sqrt{1+y'^2}}{\sqrt{2g(-y)}}$. We want to find the curve that minimizes T (Brachistochrone = ancient Greek for "shortest time").

- (a) Use Lagrange's equations to write down a differential equation for the solution.
- (b) Solve the differential equation and adapt the constants such that the boundary conditions are fulfilled.
- (c) Compare with the results of problem 1 of Blatt 1.

Remark: This problem was first posed and solved in Basel in 1696 by Johann Bernoulli!