

## Mechanik, Herbstsemester 2022

### Blatt 10

Abgabe: 29.11.2022, 12:00H, **entweder auf adam in den entsprechenden Ordner, oder in das Fach im Treppenhaus 4. Stock!**

Tutor: Julian Arnold Zi.: 4.10; julian.arnold@unibas.ch

(1) **Canonical transformations** (5 Punkte)

We consider a system with one degree of freedom ( $f = 1$ ) and want to study canonical transformations from  $(q, p)$  to  $(Q, P)$  generated by  $F_2(q, P, t)$  which leads to

$$Q = \frac{\partial F_2}{\partial P} \quad p = \frac{\partial F_2}{\partial q}.$$

Construct suitable generators  $F_2$  for the following cases:

- (a) Identical transformation:  $Q = q, P = p$ .
- (b) Galilei transformation, for which  $Q, P$  belong to a system moving with velocity  $v$  with respect to the original system:  $q = Q + vt, p = P + mv$ . Write down the new Hamiltonian  $K(Q, P)$  that is related to the original Hamiltonian  $H$  by the relation  $K = H + \frac{\partial F_2}{\partial t}$ .
- (c) A general point transformation from  $q$  to the new coordinate  $Q$ , i.e.,  $Q = g(q, t)$ . Construct a generator  $F_2$  such that the new momentum  $P$  is *proportional* to the original momentum  $p$ . Show that the transformation rule for the momentum can also be obtained from the fact that the Lagrangian is invariant under a point transformation:  $L(q, \dot{q}; t) = \tilde{L}(Q, \dot{Q}; t)$ .

(2) **Phase portraits and Liouville's theorem** (5 Punkte)

Liouville's theorem states that the volume in phase space occupied by a collection of systems remains constant over time.

Consider the dimensionless Hamiltonian  $H(q, p) = \frac{p^2}{2} + V(q)$ , where  $V(q)$  is given by

- (a)  $V(q) = \frac{q^2}{2}$  (harmonic oscillator).
- (b)  $V(q) = \frac{q^2}{2} + q^4$  (anharmonic oscillator).
- (c)  $V(q) = 1 - \cos(q)$  (pendulum;  $q$  corresponds to the deflection angle).

For each of these examples, draw phase portraits by plotting phase-space orbits  $(q(t), p(t))$  for a number of equidistant energies. Discuss the difference between open and closed orbits in (c). What happens for the initial condition  $q = 0, p = 2$  in (c)?

Consider now the rectangular phase-space volume  $-\frac{1}{2} \leq q \leq \frac{1}{2}, 1 \leq p \leq 3$  at  $t = 0$ . Analyze its time evolution qualitatively by carefully sketching (or numerically calculating) its shape for different times (at least for a time  $t \approx 1$  and for a time  $t \gg 1$ ).